

A STABLE RECURSIVE ALGORITHM FOR EVALUATION OF ULTIMATE RUIN PROBABILITIES

BY MARC GOOVAERTS AND FLORIAN DE VYLDER*

K.U. Leuven and U.C. Louvain, Belgium

ABSTRACT

Probabilities of ruin are solutions of differential or integrodifferential equations. Solving such equations numerically can be performed by means of approximate quadrature formulae for the convolution part of the equation. In this contribution it is shown how applicable recursion formulae, giving results within a prescribed tolerance level, can be obtained. Some numerical results are displayed.

KEYWORDS

Ruin probability, numerical analysis, ordering of risks.

1. INTRODUCTION

Gerber (1982) introduced a method for approximating the distribution of aggregate claims and their corresponding stop-loss premium by means of a discrete compound Poisson distribution and its corresponding stop-loss premium. This discretization is an important step in the numerical evaluation of the distribution of aggregate claims, because recent results on recurrence relations for probabilities by PANJER (1981) and SUNDT and JEWELL (1981) only apply to discrete distributions. The discretization technique is efficient in a certain sense, because a properly chosen discretization gives raise to numerical upper and lower bounds on the stop-loss premium, giving the possibility of calculating the numerically estimates for the error on the final numerical results. For calculating the infinite time ruin probability numerically one has to solve the following integral equation, according to GERBER (1979):

$$(1) \quad \psi(x) = \frac{\lambda}{c} \int_0^x \psi(x-y)(1-F(y)) dy + \frac{\lambda}{c} \int_x^\infty (1-F(y)) dy$$

with

$$(2) \quad \psi(0) = \frac{\lambda p}{c}, \quad c = \lambda p(1 + \eta)$$

and where

$$(3) \quad p = \int_0^\infty (1-F(y)) dy.$$

* We would like to thank the referees for some remarks on an earlier draft of the paper

In the sequel we write $h(x)$ for

$$(4) \quad h(x) = \int_x^{\infty} (1 - F(y)) dy$$

such that $h(0) = p$.

In fact, for solving equation (1), use could be made of a discretization technique for approximating the integrals in the equation in order to get a system of linear equations in the unknown probabilities $\psi(x_1), \psi(x_2), \dots, \psi(x_n)$. However, not every discretization is effective to get numerically stable results, on the contrary the propagation of the error induced by the discretization technique provides us, in general, with unstable numerical results. In this contribution we will derive an efficient discretization technique which allows us to calculate $\psi(x)$ numerically within a given tolerance by means of a stable recursive algorithm. As by-product of the method an estimate for the error in the numerical result is obtained.

2. AN EFFICIENT DISCRETIZATION FOR THE CONVOLUTION INTEGRAL

In the sequel we define the convolution product by

$$\varphi * H(x) = \int_0^x \varphi(x-y) dH(y)$$

where the integral is taken over the closed interval. The iterative procedure for calculating numerically the infinite time ruin probabilities is based on the following result:

THEOREM. *The function ψ (denoting the infinite time ruin probability) is a solution of the equation*

$$(5) \quad \psi * (p - h) = p(1 + \eta)\psi - h.$$

In that equation or in any equivalent equation replace (h, p) by (h_0, p_0) and let ψ_0 be the corresponding solution. Then $\psi \leq \psi_0$ ($\psi \geq \psi_0$) if h_0 is decreasing $h_0 \geq 0$, and $h/p \leq h_0/p_0$ ($h/p \geq h_0/p_0$).

PROOF. See appendix.

COROLLARY. *In case $p = p_0$ the inequality condition of the theorem reduces to $h \leq h_0$ ($h \geq h_0$).*

REMARKS. For the proof of the theorem to hold it is not necessary that h_0 can be written as $h_0(x) = \int_x^{\infty} (1 - F_0(x)) dx$ where F_0 is a distribution function.

In this extended version of the theorem, we do not have the additional assumption $h_0(0) = p$. Indeed, $h(0) = p$, so in order to get an upper bound we have to suppose $p \leq h_0(0)$ and in order to get a lower bound we have to suppose $p \geq h_0(0)$.

The theorem, or another version of it, has given us a possibility to obtain analytical upper and lower bounds on infinite time ruin probabilities in case of constraints on claim size distribution, as explained in GOOVAERTS and DE VYLDER (1983). Now it will enable us to deduce numerical bounds on infinite time ruin probabilities because an application of the theorem will provide us with a stable recursive algorithm. Our aim consists in calculating $\psi(x)$. In order to obtain numerical upper and lower bounds (to obtain an error estimate) the following procedure is applied. The underlying motivation of it follows directly from an inspection of figure 1 and an application of the theorem.




FIGURE 1

Practical Procedure

(i) Consider the following subdivision of the interval $[0, x]$

$$\left[0, \frac{x}{n}\right], \left[\frac{x}{n}, 2\frac{x}{n}\right], \dots, \left[\frac{n-1}{n}x, x\right].$$

(ii) As indicated in fig. 1 we consider

$$\forall y \in \left[\frac{i}{n}x, \frac{i+1}{n}x\right] \quad h_u(y) = h\left(\frac{i}{n}x\right)$$

$$h_l(y) = h\left(\frac{(i+1)x}{n}\right).$$

Consequently

$$h(y) \leq h_u(y) \quad \forall y \geq 0$$

$$h(y) \geq h_l(y) \quad \forall y \geq 0.$$

(iii) Let $\psi_u(x)$ be the solution of

$$\psi_u(k) \left(1 - \frac{\lambda^p 1}{c}\right) = \frac{\lambda}{c} \left(1 - \frac{\lambda^p 1}{c}\right) h_u(k) - \frac{\lambda}{c} \int_0^k \psi'_u(k-y) h_u(y) dy.$$

Because $h_u(y)$ is a piecewise constant function, the integral appearing in the r.h.s. can be worked out as follows: let $k = j(x/n)$, then:

$$\begin{aligned} \psi_u\left(j\frac{x}{n}\right) \left(1 - \frac{\lambda^p 1}{c}\right) &= \frac{\lambda}{c} \left(1 - \frac{\lambda^p 1}{c}\right) h\left(j\frac{x}{n}\right) \\ &\quad - \frac{\lambda}{c} \sum_{i=0}^{j-1} h\left(i\frac{x}{n}\right) \int_{i(x/n)}^{(i+1)(x/n)} \psi'_u\left(j\frac{x}{n} - y\right) dy. \end{aligned}$$

This equation can be cast into the form:

$$(6) \quad \psi_u\left(j\frac{x}{n}\right) = \frac{\lambda}{c} h\left(j\frac{x}{n}\right) + \frac{\lambda}{c} \sum_{i=1}^j \left[h\left((i-1)\frac{x}{n}\right) - h\left(i\frac{x}{n}\right) \right] \psi_u\left((j-i)\frac{x}{n}\right).$$

We also get, proceeding along the same lines,

$$(7) \quad \begin{aligned} \psi_l\left(j\frac{x}{n}\right) &= \frac{\lambda}{c} \left[1 - \frac{\lambda}{c} (p_1 - h\left(\frac{x}{n}\right)) \right]^{-1} \left[h\left(j\frac{x}{n}\right) \right. \\ &\quad \left. + \sum_{i=1}^{j-1} \left[h\left(i\frac{x}{n}\right) - h\left((i+1)\frac{x}{n}\right) \right] \psi_l\left((j-i)\frac{x}{n}\right) \right] \end{aligned}$$

with of course

$$(8) \quad \psi_u(0) = \psi_l(0) = \frac{\lambda p}{c}.$$

Starting from (8) we calculate recursively by means of (6) and (7) the couple $(\psi_u(j(x/n)), \psi_l(j(x/n)))$ for $j = 1, 2, \dots, n$ to obtain two approximations to $\psi(x)$, namely $\psi_u^{(n)}(x)$, $\psi_l^{(n)}(x)$ where we added explicitly the index n to denote the dependence on n . The following inequalities are obtained

$$(9) \quad \psi_l^{(n)}(x) \leq \psi(x) \leq \psi_u^{(n)}(x).$$

Also from the result of the above quoted theorem we conclude that

$$\psi_l^{(2^n)}(x) \text{ is not decreasing in } n$$

$$\psi_u^{(2^n)}(x) \text{ is not increasing in } n$$

Hence $\psi(x)$ can be approximated by

$$\psi(x) \approx \frac{1}{2} \psi_l^{(n)}(x) + \frac{1}{2} \psi_u^{(n)}(x)$$

with an upper bound for the error given by

$$\psi_u^{(n)}(x) - \psi_l^{(n)}(x).$$

In order to obtain a result within a prescribed tolerance level ε , n is chosen large enough such that

$$\psi_u^{(n)}(x) - \psi_l^{(n)}(n) < \varepsilon$$

and $n = 2^k$ (k integer).

In order to examine the stability of the numerical procedure, in fact, in order to examine the propagation of errors induced by this recursive algorithm, we suppose that $\psi(x/n), \dots, \psi((j-1)x/n)$ are calculated with an error $\varepsilon_1, \dots, \varepsilon_{j-1}$. Let $\varepsilon = \max\{\varepsilon_1, \dots, \varepsilon_{j-1}\}$ then the error ε_j on $\psi(j(x/n))$ is bounded by:

$$\varepsilon_j < \frac{\lambda}{c} \sum_{i=1}^j \left(h\left((i-1)\frac{x}{n}\right) - h\left(i\frac{x}{n}\right) \right) \quad \varepsilon_j < \varepsilon.$$

Consequently there is no cumulative effect of propagation of errors. Let us compare the kind of recursion relation with the recursive algorithm of PANJER (1981) and SUNDT and JEWELL (1981) for the calculation of the distribution function of a compound Poisson variable, where

$$g_j = \frac{\lambda}{j} \sum_{i=1}^j i f_i g_{j-i}.$$

In case λ is relatively small no problem arises as far as the propagation of errors is concerned.

In case λ is relatively large however (and this is exactly the case where it is interesting to apply such kind of a scheme) the recursive algorithm is unfortunately not very stable as far as the propagation of errors is concerned. Indeed let $\varepsilon_1, \dots, \varepsilon_{j-1}, \varepsilon_j$ denote the errors on g_1, \dots, g_j respectively. In case $\varepsilon = \max\{\varepsilon_1, \dots, \varepsilon_{j-1}\}$ then

$$\varepsilon_j \leq \frac{\lambda}{j} \sum_{i=1}^j i f_i \quad \varepsilon \approx \mu \frac{\lambda}{j} \cdot \varepsilon.$$

Consequently as long as $j < \lambda$ the upper bound of the error behaves like

$$\varepsilon_j \approx \lambda^j \cdot c$$

which of course can cause a lot of unexpected difficulties in actual application of the recursion algorithm.

3. ILLUSTRATION OF THE METHOD

In VAN WOUWE, DE VYLDER and GOOVAERTS (1982) the present results are successfully applied to the numerical calculation of bounds on infinite time ruin probabilities in case of constraints on claim size distributions. Use has been made of some of the analytical upper and lower bounds on stop-loss premiums.

Let us still remark that the rate of convergence of the recursive algorithm is determined by the rate of convergence to zero of the ruin probability when $x \rightarrow \infty$. Therefore we have selected an application which from the numerical point of view has a relatively low speed of convergence.

We consider the case of Pareto claims, $F_X(x) = 1 - 1/(1+x)^2$, $p = 1$, $\eta = 0.2$ and of course $h(x) = 1/(1+x)$. The following results are obtained.

Upper Bound

$n \backslash x$	10	50	100
20	0.455529	0.193577	0.119406
40	0.449979	0.164704	0.087263
80	0.439944	0.153144	0.076432
160	0.439944	0.148211	0.072358

Lower Bound

$n \backslash x$	10	50	100
20	0.411083	0.121643	0.058221
40	0.422112	0.129821	0.061631
80	0.428309	0.135709	0.064429
160	0.431619	0.139413	0.066421

APPENDIX PROOF OF THE THEOREM

Use will be made of the following result well known from renewal theory:

LEMMA. *If H is a strictly defective distribution function and if f is bounded, then the renewal equation*

$$(A1) \quad \xi = f + \xi * H$$

has a unique bounded solution ξ . If $f \geq 0$, then $\xi \geq 0$. If $f \leq 0$, then $\xi \leq 0$.

By means of one partial integration the equation (1) can be cast into the form

$$(A2) \quad \psi * (p - h) = p(1 + \eta)\psi - h.$$

This relation can still be displayed as:

$$(A3) \quad p - p(1 + \eta)\psi = (1 - \psi) * (p - h).$$

By the definition of ψ_0 , we still have

$$(A4) \quad p_0 - p_0(1 + \eta)\psi_0 = (1 - \psi_0) * (p_0 - h_0).$$

From (A3) and (A4) we deduce

$$(A5) (1 + \eta)(\psi - \psi_0) = h/p - h_0/p_0 + (\psi - \psi_0) * (1 - h/p) + \psi_0 * (h_0/p_0 - h/p).$$

Then the above mentioned lemma can be applied with

$$\xi = \psi - \psi_0$$

$$H = (1 - h/p)/(1 + \eta)$$

$$(A6) \quad f = (1 - \psi_0) * (h/p - h_0/p_0)/(1 + \eta).$$

Then of course f is bounded. Moreover because $h/p \geq h_0/p_0$ we have $f \geq 0$. The function H is a distribution function (it is increasing and $H \geq 0$), in fact it is a strictly defective distribution function because $H(\infty) = 1/(1 + \eta)$. Hence the result of the theorem follows.

REFERENCES

- GERBER, H. U. (1979) *An Introduction to Mathematical Risk Theory* Homewood Richard D. Irwin, Inc.
- GERBER, H. U. (1982) On the Numerical Evaluation of the Distribution of Aggregate Claims and its Stop-Loss Premiums. *Insurance, Mathematics and Economics*, **1**, 13-18.
- GOOVAERTS, M. and DE VYLDER, F. (1983) Upper and Lower Bounds on Infinite Time Ruin Probabilities in Case of Constraints on Claim Size Distributions *Journal of Econometrics*, **23**, 77-90.
- PANJER, H. H. (1981) Recursive Evaluation of a Family of Compound Distributions. *Astin Bulletin* **12**, 22-26.
- SUNDT, B. and JEWELL, W. S. (1981) Further Results on Recursive Evaluation of Compound Distributions *Astin Bulletin* **12**, 27-39.
- WOUWE, M. VAN, DE VYLDER, F. and GOOVAERTS, M. (1982) Numerical Bounds on Infinite Time Ruin Probabilities in Case of Constraints on Claim Size Distributions. (To be presented at the 22nd International Congress of Actuaries, Sydney 1984).

