

RECURSIVE EVALUATION OF A FAMILY
OF COMPOUND DISTRIBUTIONS*

HARRY H. PANJER

University of Waterloo, Ontario, Canada

1 INTRODUCTION

Compound distributions such as the compound Poisson and the compound negative binomial are used extensively in the theory of risk to model the distribution of the total claims incurred in a fixed period of time. The usual method of evaluating the distribution function requires the computation of many convolutions of the conditional distribution of the amount of a claim given that a claim has occurred. When the expected number of claims is large, the computation can become unwieldy even with modern large scale electronic computers.

In this paper, a recursive definition of the distribution of total claims is developed for a family of claim number distributions and arbitrary claim amount distributions. When the claim amount is discrete, the recursive definition can be used to compute the distribution of total claims without the use of convolutions. This can reduce the number of required computations by several orders of magnitude for sufficiently large portfolios.

Results for some specific distributions have been previously obtained using generating functions and Laplace transforms (see PANJER (1980) including discussion). The simple algebraic proof of this paper yields all the previous results as special cases.

2. THE FAMILY OF CLAIM NUMBER DISTRIBUTIONS

Consider the family of claim number distributions satisfying the recursion

$$(1) \quad p_n = p_{n-1}(a + b/n), \quad n = 1, 2, 3, \dots$$

where p_n denotes the probability that exactly n claims occur in the fixed time interval. Members of this family are

a) Poisson distribution:

$$1. \quad p_n = \frac{e^{-\lambda} \lambda^n}{n!}, \quad n = 0, 1, 2, \dots$$

* The author is grateful to the referee for pointing out an error in the original draft. This research was supported by the Natural Sciences and Engineering Research Council of Canada.

2. $p_n/p_{n-1} = \lambda/n, \quad p_0 = e^{-\lambda}$
3. $a = 0, \quad b = \lambda$

b) Binomial distribution

1. $p_n = \binom{N}{n} p^n (1-p)^{N-n}, \quad n = 0, 1, 2, \dots, N$
 $p_n = 0, \quad n = N+1, N+2, \dots$
2. $p_n/p_{n-1} = (N-n+1)p/(n(1-p)), \quad p_0 = (1-p)^N$
3. $a = -p/(1-p), \quad b = (N+1)p/(1-p)$

c) Negative binomial distribution

1. $p_n = \binom{\alpha+n-1}{n} p^n (1-p)^\alpha, \quad n = 0, 1, 2, \dots$
2. $p_n/p_{n-1} = (\alpha+n-1)p/n, \quad p_0 = (1-p)^\alpha$
3. $a = p, \quad b = (\alpha-1)p$

d) Geometric distribution (Negative binomial with $\alpha = 1$):

1. $p_n = (1-p)p^n, \quad n = 0, 1, 2, \dots$
2. $p_n/p_{n-1} = p, \quad p_0 = 1-p$
3. $a = p, \quad b = 0$

SUNDI and JEWELL (1981) show that these are the only members of this family.

3 THE RECURSIVE FORMULA

Consider the compound distribution with distribution function

$$(2) \quad G(x) = \sum_{n=1}^{\infty} p_n F^{*n}(x), \quad x > 0$$

$$= p_0, \quad x = 0$$

for arbitrary claim amount distribution $F(x), x > 0$. For notational convenience, assume that $F(x), x > 0$, is of the continuous form. Corresponding results will be given for the discrete case. Then the density of total claims is

$$(3) \quad g(x) = \sum_{n=1}^{\infty} p_n f^{*n}(x), \quad x > 0$$

$$= p_0, \quad x = 0$$

when $f(x)$ is the density associated with $F(x)$

In the theorem which follows, the following two relations will be used:

$$(I) \quad \int_0^x f(y) f^{*n}(x-y) dy = f^{*(n+1)}(x), \quad n = 1, 2, 3, \dots$$

$$(II) \quad \int_0^x y f(y) f^{*n}(x-y) / f^{*(n+1)}(x) dy = x/(n+1), \quad n = 1, 2, 3, \dots$$

Relation (I) is the usual recursive definition of convolutions. The left side of relation (II) is the conditional mean of any element of a sum consisting of $n+1$ independent and identically distributed elements, given that the sum is exactly x . The mean is $x/(n+1)$ as a result of the symmetry in the elements of the sum. Relation (II) is used in Buhlmann and Gerber's discussion to PANJER (1980) to develop an alternate proof of the result described in that paper.

Theorem

For p_n and $g(x)$ defined by (1) and (3) respectively, and $f(x)$ any distribution of the continuous type for $x > 0$, the following recursion holds.

$$(4) \quad g(x) = p_1 f(x) + \int_0^x (a + by/x) f(y) g(x-y) dy, \quad x > 0.$$

Proof

Substituting (3) into the right side of (4) results in

$$\begin{aligned} p_1 f(x) + \int_0^x (a + by/x) f(y) g(x-y) dy &= p_1 f(x) + \int_0^x (a + by/x) f(y) \\ &\quad \sum_{n=1}^{\infty} p_n f^{*n}(x-y) dy \\ &= p_1 f(x) + \sum_{n=1}^{\infty} p_n \int_0^x (a + by/x) f(y) f^{*n}(x-y) dy \\ &= p_1 f(x) + \sum_{n=1}^{\infty} p_n \{a + b/(n+1)\} f^{*(n+1)}(x) \\ &\quad \text{(from (I) and (II))} \\ &= p_1 f(x) + \sum_{n=1}^{\infty} p_{n+1} f^{*(n+1)}(x) \quad \text{(from (1))} \\ &= p_1 f(x) + \sum_{n=1}^{\infty} p_n f^{*n}(x) \\ &= \sum_{n=1}^{\infty} p_n f^{*n}(x) \quad \text{(since } f^{*1}(x) = f(x)) \\ &= g(x), \quad \text{the required result.} \quad \text{Q.E.D.} \end{aligned}$$

If the claim amount distribution is discrete and defined on the positive integers, the corresponding recursive definition of total claims is

$$(5) \quad g_i = \sum_{j=1}^i (a + bj/i) f_j g_{i-j}, \quad i = 1, 2, 3, \dots$$

with $g_0 = p_0$; whereas the usual form is

$$(6) \quad g_i = \sum_{n=0}^i p_n f_i^n, \quad i = 0, 1, 2, 3, \dots$$

The number of computations required to obtain g_i is of order i^2 for formula (6) and of order i for formula (5). Hence, for large values of i , the reduction in computations is dramatic.

4. RESULTS FOR SPECIAL CASES

The recursive definitions of the density of total claims for the four distributions considered in Section 2 are given below.

Continuous Model

Discrete Model

$g(x)$

g_i

a) Poisson

$$\lambda e^{-\lambda} f(x) + \frac{\lambda}{x} \int_0^x y f(y) g(x-y) dy \quad \frac{\lambda}{i} \sum_{j=1}^i j f_j g_{i-j}$$

b) Binomial

$$\frac{p}{1-p} \left[N(1-p)^N f(x) + \int_0^x \{(N+1) y/x - 1\} f(y) g(x-y) dy \right]$$

$$\frac{p}{1-p} \sum_{j=1}^i \{(N+1) j/i - 1\} f_j g_{i-j}$$

c) Negative binomial

$$p \left[\alpha(1-p)^\alpha f(x) + \int_0^x \{1 + (\alpha-1) y/x\} f(y) g(x-y) dy \right]$$

$$p \sum_{j=1}^i \{1 + (\alpha-1) j/i\} f_j g_{i-j}$$

d) Geometric

$$p[(1-p) f(x) + \int_0^x f(y) g(x-y) dy] \quad p \sum_{j=1}^i f_j g_{i-j}$$

The recursion for the compound Poisson distribution with discrete claim amount distribution was originally given by ADELSON (1966) in an inventory problem. He used generating functions to obtain his result.

SUNDT and JEWELL (1981) generalize the results of the present paper. They obtain results for a more general recursion than (1) and obtain results for the case of possibly negative claims.

REFERENCES

- ADELSON, R M (1966) Compound Poisson Distributions, *Operations Research Quarterly*, 17, 73-75
- PANJER, H H. (1980) The Aggregate Claims Distribution and Stop-Loss Reinsurance, *Trans of the Society of Actuaries*, XXXII.
- SUNDT, B and JEWELL, W S (1981) Further Results on Recursive Evaluation of Compound Distributions *Astin Bulletin* 12